

p -Adic Arithmetic in SAGE

David Roe

Department of Mathematics
Harvard University

Sage Days 5

Outline

- 1 **Motivating Goals**
 - Mission
 - Applications and Objects of Interest
- 2 Mathematical tools needed
- 3 Current Status and Priorities

Outline

- 1 Motivating Goals
 - Mission
 - Applications and Objects of Interest
- 2 Mathematical tools needed
- 3 Current Status and Priorities

Outline

- 1 Motivating Goals
 - Mission
 - Applications and Objects of Interest
- 2 Mathematical tools needed
- 3 Current Status and Priorities

Why we want p -adics in Sage

To support the advancement of mathematical knowledge by providing the facility to compute with mathematically interesting objects relying on \mathbb{Q}_p .

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Mathematical Objects

We want to compute with many mathematical objects that rely on \mathbb{Q}_p :

- Spaces of p -adic modular forms and overconvergent modular forms.
- p -adic and ℓ -adic Galois representations as part of a more general framework for Galois representations.
- p -adic cohomology theories (crystalline, étale, Monsky-Washnitzer) as part of a more general framework for cohomology in Sage.
- p -adic L-functions, zeta functions, etc.
- p -adic analogues of trace-formulae.
- ???

Applications

Similarly, we can apply p -adics to a host of problems . . .

- p -adic heights for points on elliptic curves.
- Applications of p -adic differential equations.
- Applications of p -adics to quadratic forms (e.g. the Hasse principle)
- Applications of p -adics to linear algebra over global fields (e.g. Dixon's algorithm)
- ???

Applications

Similarly, we can apply p -adics to a host of problems . . .

- p -adic heights for points on elliptic curves.
- Applications of p -adic differential equations.
- Applications of p -adics to quadratic forms (e.g. the Hasse principle)
- Applications of p -adics to linear algebra over global fields (e.g. Dixon's algorithm)
- ???

Applications

Similarly, we can apply p -adics to a host of problems . . .

- p -adic heights for points on elliptic curves.
- Applications of p -adic differential equations.
- Applications of p -adics to quadratic forms (e.g. the Hasse principle)
- Applications of p -adics to linear algebra over global fields (e.g. Dixon's algorithm)
- ???

Applications

Similarly, we can apply p -adics to a host of problems . . .

- p -adic heights for points on elliptic curves.
- Applications of p -adic differential equations.
- Applications of p -adics to quadratic forms (e.g. the Hasse principle)
- Applications of p -adics to linear algebra over global fields (e.g. Dixon's algorithm)
- ???

Applications

Similarly, we can apply p -adics to a host of problems . . .

- p -adic heights for points on elliptic curves.
- Applications of p -adic differential equations.
- Applications of p -adics to quadratic forms (e.g. the Hasse principle)
- Applications of p -adics to linear algebra over global fields (e.g. Dixon's algorithm)
- ???

Applications

Similarly, we can apply p -adics to a host of problems . . .

- p -adic heights for points on elliptic curves.
- Applications of p -adic differential equations.
- Applications of p -adics to quadratic forms (e.g. the Hasse principle)
- Applications of p -adics to linear algebra over global fields (e.g. Dixon's algorithm)
- ???

Extensions

Currently, Sage does not support extensions of \mathbb{Q}_p . This will change soon.

- There will be special types for unramified extensions of \mathbb{Q}_p , eisenstein extensions of \mathbb{Q}_p , general absolute extensions and general relative extensions
- A general extension will be converted transparently into a two step extension (first an unramified extension, then an eisenstein extension)
- Krasner's lemma will be used to determine if the defining polynomial uniquely defines an extension.
- Eventually we will need a version of round 4 to split an arbitrary extension into unramified and totally ramified parts.

More on Extensions

- Underlying arithmetic will be done by the NTL classes `ZZ_pE` and `ZZ_pEX`.
- As with the base classes, different elements of the same ring or field can have different precision, necessitating casting during arithmetic.
- Elements of a general extension will cache their absolute and relative forms.

Completions

- It should be trivial in Sage to obtain the completion of a number field K at a given prime/place.
- Extensions need to happen first.
- There should be a type, “Completion of number field” that includes a local field and a map from the number field.

Polynomials

- There should be special classes to take advantage of fast NTL code.
- How should precisions of coefficients be restricted?
- One can write good algorithms in the mixed precision case.
- Need to support operations such as resultants that require linear algebra.

Linear algebra and modules

- We need to perform the standard linear algebra operations (determinants and traces, kernels, characteristic and minimal polynomials, etc). But these operations are harder in the p -adic case because of precision issues.
- As with polynomials, one can try to find an answer and find the precision separately (for some problems at least).
- As with polynomials, there is a question of how precision varies across a matrix, within a module or vector space.
- Algorithms need to be numerically stable.

Power series

- Much of the work on p -adics generalizes to power series.
- With power series over p -adics one can impose more complicated precision restrictions.
- It's probably fast enough to keep implementing power series as a Sage polynomial and a precision.
- There is some demand for bidirectional power series, with conditions on the norms of the coefficients.

\mathbb{Q}_p and \mathbb{Z}_p

- The base classes are generally in good shape, though the lazy classes need work.
- Almost all the p -adic code suffers from lack of doctests.
- The current code for the base classes is quite fast, comparable to Magma in arithmetic tests.

Priorities

What is our plan for p -adics in Sage?